



GCCHD

Greene County Combined Health District
360 Wilson Drive
Xenia, Ohio 45385

News Release

For Immediate Release
June 22, 2010

Contact: Sara T. Pappa, Public Information Officer
(937) 374-5600 or 1-866-858-3588

Pertussis “Whooping Cough” Cases Reported in Greene County

(Xenia, OH) -- The Greene County Combined Health District (GCCHD) has seen an increase in the number of cases of “Whooping Cough”, medically known as Pertussis, over the past week with 7 cases reported in the Beavercreek area.

Pertussis (whooping cough) is a very contagious disease caused by a type of bacteria called *Bordetella pertussis*. Among vaccine-preventable diseases, pertussis is one of the most commonly occurring ones in the United States.

There is high vaccine coverage for children nationwide. However, protection from the childhood vaccine fades over time. Adolescents and adults need to be revaccinated, even if they were completely vaccinated as children.

Also, pertussis vaccines are very effective but not 100% effective. A fully vaccinated person can catch this very contagious disease. When you or your child develops a cold that includes a prolonged or severe cough, it may be pertussis. The best way to know is to contact your doctor.

Pertussis can cause serious illness in infants, children and adults. The disease starts like the common cold, with runny nose or congestion, sneezing, and maybe mild cough or fever. But after 1–2 weeks, severe coughing begins. Infants and children with the disease cough violently and rapidly, over and over, until the air is gone from their lungs and they're forced to inhale with a loud "whooping" sound. Pertussis is most severe for babies; more than half of infants less than 1 year of age who get the disease must be hospitalized. About 1 in 20 infants with pertussis get pneumonia (lung infection), and about 1 in 100 will have convulsions. In rare cases, pertussis can be deadly, especially in infants.

People with pertussis usually spread the disease by coughing or sneezing while in close contact with others, who then breathe in the pertussis bacteria. Many infants who get pertussis are infected by parents, older siblings, or other caregivers who might not even know they have the disease.

Since the 1980s, there's been an increase in the number of cases of pertussis, especially among teens (10–19 years of age) and babies less than 6 months of age. In 2008 there were more than 13,000 reported cases including 18 deaths from pertussis nationally.

The best way to prevent pertussis is to get vaccinated. In the US, the recommended pertussis vaccine for children is called DTaP. This is a safe and effective combination vaccine that protects children against three diseases: diphtheria, tetanus, and pertussis. For maximum protection against pertussis, children need five DTaP shots. The first three shots are given at 2, 4, and 6 months of age. The fourth shot is given between 15 and 18 months of age, and a fifth shot is given when a child enters school, at 4–6 years of age. Parents can also help protect infants by keeping them away as much as possible from anyone who has cold symptoms or is coughing.

Vaccine protection for pertussis, tetanus, and diphtheria can fade with time. Before 2005, the only booster vaccine available contained tetanus and diphtheria (called Td), and was recommended for adolescents and adults every 10 years. Today there are boosters for adolescents and adults that contain tetanus, diphtheria, *and* pertussis (called Tdap). Pre-teens going to the doctor for their regular check-up at age 11 or 12 years should get a dose of Tdap. Adults who didn't get Tdap as a pre-teen or teen should get one dose of Tdap instead of the Td booster. Most pregnant women who were not previously vaccinated with Tdap should get one dose of Tdap postpartum before leaving the hospital or birthing center. Getting vaccinated with Tdap is especially important for families with and caregivers of new infants.

The easiest thing for adults to do is to get Tdap instead of their next regular tetanus booster—that Td shot that they were supposed to get every 10 years. The dose of Tdap can be given earlier than the 10-year mark, so it's a good idea for adults to talk to a healthcare provider about what's best for their specific situation.

For more information, please call Amy Schmitt at (937) 374-5638 or Don Brannen at (937) 374-5660 at the Greene County Combined Health District or visit the Centers for Disease Control and Prevention at www.cdc.gov.

**Greene County Combined Health District – Your center for public health services
and health information in Greene County for over 90 years.**

Healthy People...Healthy Communities

360 Wilson Drive

Xenia, OH 45385

937-374-5600 * 866-858-3588 * Fax 937-374-5675

www.gcchd.org